



Coding-Complete Genome Sequence of SARS-CoV-2 Isolate from Bangladesh by Sanger Sequencing

M. Moniruzzaman,^a  Mohammad Uzzal Hossain,^b M. Nazrul Islam,^c M. Hadisur Rahman,^a Irfan Ahmed,^a Tahia Anan Rahman,^a Arittra Bhattacharjee,^{b,d} M. Ruhul Amin,^e Asif Rashed,^f Chaman Ara Keya,^d Keshob Chandra Das,^a  M. Salimullah^a

^aMolecular Biotechnology Division, National Institute of Biotechnology, Dhaka, Bangladesh

^bBioinformatics Division, National Institute of Biotechnology, Savar, Dhaka, Bangladesh

^cPlant Biotechnology Division, National Institute of Biotechnology, Savar, Dhaka, Bangladesh

^dDepartment of Biochemistry and Microbiology, North South University, Bashundhara, Dhaka, Bangladesh

^eCenter for Medical Biotechnology, MIS, Directorate General of Health Services, Dhaka, Bangladesh

^fDepartment of Microbiology, Mugda Medical College, Dhaka, Bangladesh

ABSTRACT A coding-complete genome sequence of a severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) isolate was revealed. The sample for the virus was isolated from a female patient from Dhaka, Bangladesh, suffering from coronavirus disease-2019 (COVID-19).

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), a member of the *Coronaviridae* family and *Betacoronavirus* genus, is the causative agent of pandemic coronavirus disease-2019 (COVID-19). In Bangladesh, the rate of positive cases and the death toll from COVID-19 are increasing at an alarming rate (<https://corona.gov.bd/>). To understand the genomic characteristics of SARS-CoV-2 in Bangladesh, several isolates have been sequenced and deposited in GISAID (<https://www.gisaid.org/>). However, those isolates have been sequenced using a next-generation sequencing platform, except for the one we are reporting. In this study, we sequenced the viral genome by Sanger sequencing technology, which is a gold standard method and is necessary for thorough genomic analysis (1).

The isolate (SARS-CoV-2/human/BGD/NIB_01/2020) was collected from an oropharyngeal specimen on 11 May 2020. The patient was a 28-year-old saleswoman who tested positive (via reverse transcriptase PCR [RT-PCR]) for COVID-19 with symptoms of cough, mild fever, and throat congestion. (all applicable international, national, and/or institutional guidelines for the care and use of animals were followed; ethical approval number NIBREC2020-01). The viral RNA was extracted directly from the patient's specimen using the PureLink viral RNA/DNA minikit (Invitrogen). The viral RNA was then converted into cDNA using a SuperScript VILO cDNA synthesis kit (Invitrogen).

To cover the whole genome of the virus, 48 pairs of primers were designed by following two conditions: (i) their sequence is conserved among all the available SARS-CoV-2 isolates, and (ii) the terminal of the amplicons will overlap the adjacent amplicon (Table 1). These primers underwent PCR and generated 96 amplicons, which were visualized using 1.5% agarose gel electrophoresis. The PCR products were then purified using the PureLink PCR purification kit (Thermo Fisher Scientific, USA). These purified amplicons were finally sequenced with 2× coverage using the Sanger dideoxy method by "ABI 3500" with a BigDye Terminator version 3.1 cycle sequencing kit (Applied Biosystems, USA).

The raw reads were assembled using DNA Sequence Assembler version 4 (2013)

Citation Moniruzzaman M, Hossain MU, Islam MN, Rahman MH, Ahmed I, Rahman TA, Bhattacharjee A, Amin MR, Rashed A, Keya CA, Das KC, Salimullah M. 2020. Coding-complete genome sequence of SARS-CoV-2 isolate from Bangladesh by Sanger sequencing. *Microbiol Resour Announc* 9:e00626-20. <https://doi.org/10.1128/MRA.00626-20>.

Editor John J. Dennehy, Queens College

Copyright © 2020 Moniruzzaman et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to M. Salimullah, salim2969@gmail.com.

Received 29 May 2020

Accepted 19 June 2020

Published 9 July 2020

TABLE 1 Information about primers, amplicon size, and overlapping length

Amplicon	Primer	Sequence	Product size (bp)	Overlapping length (bp)
1	Forward	AGGTTTATACCTCCCAGG	765	
1	Reverse	CACCACTGCTATGTTAGTG		131
2	Forward	CGCAAGGTTCTTCGTA	797	
2	Reverse	AGACTATGCTCAGGTCTAC		100
3	Forward	AAGAAGGTGCCACTACTTG	794	
3	Reverse	GTTAGTTAGCCACTGCGAA		
4	Forward	ACTGAGACTCATTGATGCTA	788	114
4	Reverse	TCACACTCTTGTAAACCTTG		
5	Forward	AGAAAAGTACTGTGCCCTG	783	130
5	Reverse	ACCACTGTTGGTTTACCTT		
6	Forward	GATGGAACCTAACCCAGTTG		131
6	Reverse	GTCACTAACAAGAGTGGCAG	781	
7	Forward	GAAGAAGTTACAACAACCTCTGG	718	110
7	Reverse	AAACTGTAGCTGGCACTTTG		
8	Forward	TGTAGCGTCACTTACAACA	746	136
8	Reverse	CAGTGGCAAGATAACAGTTG		
9	Forward	CTCTACGTGTTGAGGCTTT	720	158
9	Reverse	CATCGTAATAGGACCTTGT		
10	Forward	TTGTGCTAGTGAGTACACTG	760	130
10	Reverse	AATGTCCTACAATTCTGG		
11	Forward	TGATGTAATGAGTCAGAGG	737	150
11	Reverse	AATAGCCTCTCTGTAACCAG		
12	Forward	TTCTTTAATCTACTCAACCGC	706	90
12	Reverse	CTGTAGTGACAAGTCTCTCG		
13	Forward	ATGCTAATGGAGGTAAAGGC	701	118
13	Reverse	ACAACATATGCCAGTAACCTC		
14	Forward	CTTTTATTTCAGCAGCTCGG	714	115
14	Reverse	GTGCGTAATATCGTCCA		
15	Forward	GCTGATTGACACATGGTT	812	133
15	Reverse	GGTAAGAATGAGTAAACTGGTG		
16	Forward	CCTATTGGTGCCTTGGACATA	727	196
16	Reverse	AACCTCTAACCTTACAGATG		
17	Forward	CTTGTGTCATCTGCAAAG	767	146
17	Reverse	TCGATTGAGAAACCACTGT		
18	Forward	TTGTTGACAGGCAAACAGC	770	112
18	Reverse	ACCATCATCATACACAGTTCT		
19	Forward	TGACATGGTGGATATGGTG	794	121
19	Reverse	GTTTATGTCTACAGCACCT		
20	Forward	AATTGTTGGCTCAATGTG	787	172
20	Reverse	GCAACAGGACTAACGCTATTA		
21	Forward	GGAAATCCAACAGGTTAGA	795	155
21	Reverse	ACAGGGTCATTAGCACAAGT		
22	Forward	GTTGCCACATAGATCATCAA	790	90
22	Reverse	AAACATACCAACAGCTTCGC		
23	Forward	GCAGACCTCGTCTATGCTT	813	233
23	Reverse	GCACGTAGTGCCTTATCT		
24	Forward	CCACTTCAGAGAGCTAGGTG	782	147
24	Reverse	GTGAGGGTTTCTACATCACT		
25	Forward	ATTGAAATCAATAGCCGCA	775	114
25	Reverse	ATCTGGTAAGGAAGGTACA		
26	Forward	GTCTGAAGCAAAATGTGGA	805	117
26	Reverse	GAGCTTTCACTACAGGTGTT		
27	Forward	TGTGTGCTAATGGACAAGTT	784	142
27	Reverse	TCAAAACACTCTACACGAGC		
28	Forward	CTTCTGCTCGCATAGTGTAT	769	132
28	Reverse	CAAGAGTGAGCTGTTCACT		
29	Forward	AATAGGCCTGGTAAGAGAAAT	790	191
29	Reverse	GTACATAAGTGGTATGAGGTG		
30	Forward	AGCTAGGTTTCTACAGGTG	756	139
30	Reverse	CTTTGTCACTACAAGGCTGT		
31	Forward	GTAGAAAGGTCAACACATGG	733	152
31	Reverse	ATAGAAAAGTGTACTTCACCC		
32	Forward	GCTTAGCTTGTGGTTAC	808	144
32	Reverse	CCACCTAACTGACTATGACT		139

(Continued on next page)

TABLE 1 (Continued)

Amplicon	Primer	Sequence	Product size (bp)	Overlapping length (bp)
33	Forward	CAAGAATTAAACCCAGGAG	758	
33	Reverse	GCATCAGAGACAAAGTCATT		155
34	Forward	CACATTAACATTAGCTGTACCC	781	
34	Reverse	TGACTAGAGACTAGTGGCA		182
35	Forward	AAGGGTACTGCTGTTATGT	775	
35	Reverse	TTAATAGGCGTGTGCTTAGA		116
36	Forward	TCAGCCTTTCTATGGACC	794	
36	Reverse	TCCAAGCTATAACGCAGC		104
37	Forward	TTAGAGGTGATGAAGTCAGA	760	
37	Reverse	TGTCAGCCCCATTAAACA		149
38	Forward	TAACCAGGTTGCTGTTCTT	797	
38	Reverse	CAATCATTCATCTGTGAGCA		191
39	Forward	CAGATCCATAAAACCAAGC	771	
39	Reverse	GCAAGAAGACTACACCATGA		137
40	Forward	TCAGAGCTCTGCTAATCTTG	759	
40	Reverse	GTAATTGACTCCTTGAGC		137
41	Forward	TTGCATAGTAATGGTGACA	798	
41	Reverse	AGCTGGTAATAGTCTGAAGTG		120
42	Forward	GCACAACAAGTCTTATTCT	784	
42	Reverse	CCATAACAGCCAGAGGAAAA		170
43	Forward	GCAGATTCCAACGGTACT	707	
43	Reverse	TAGTAACCTGAAAGTCAACG		117
44	Forward	GCTACAGGATTGCAACTAT	785	
44	Reverse	TTTCATGTTCGTTAGGCGT		174
45	Forward	CACTTGCTCACACTAAA	791	
45	Reverse	TCTGGACTGCTATTGGTGT		180
46	Forward	CAGATTCAACTGGCAGTAAC	793	
46	Reverse	TTTCCCTGGGTTTGTCTGG		187
47	Forward	CTGCTTGACAGATTGAACCA	698	
47	Reverse	CTTGTGCTATGTAGTTACGAGA		242
48	Forward	ATGAAACTCAAGCCTACCG		
48	Reverse	CCTTCGTGCAGGTCAATA	518	

(Heracle BioSoft) and verified with SeqMan Pro version 14.1 (DNAStar, Madison, WI). After assembly, 48 contigs with 94 overlapping regions were obtained. These overlapping regions were visualized using CLC Genomics Workbench version 20.0.4 and merged with EMBOS: merger (2).

The assembled viral genome consists of a single-stranded positive (+) RNA that is 29,724 nucleotides long. The NCBI BLASTN program (3) showed that the genome was mostly similar to SARS-CoV-2/human/BGD/CHRF_0001/2020 (GenBank accession number MT476385.1). From NCBI, the FASTA sequences of 7 mostly similar genomes from Bangladesh, India, Sri Lanka, and the United States were taken along with the reference genome. Another 16 genomes of SARS-CoV-2 that were isolated in Bangladesh were collected from GISAID (<https://www.gisaid.org/>). The genomes were aligned with MAFFT version 7 using default parameters (4). The phylogenetic tree was constructed using FastTree version 2.1.10 (5) through the Galaxy platform (6). Here, the tree was built by nucleotide alignment using the generalized time-reversible model (GTR) plus the CAT nucleotide evolution model (GTR+CAT). The tree was visualized using iTOL (7), where the tree structure was rerooted on the position of reference isolate SARS-CoV-2 Wuhan-Hu-1.

The genome has 8 nucleotide differences from the closest isolate. Interestingly, except for isolate SARS-CoV-2/human/BGD/CHRF0001/2020, the other strains of SARS-CoV-2 from Bangladesh showed separate clades and distant genetic relations. The tree also demonstrated that our viral genome and three isolates from the United States share an ancestor (Fig. 1).

Data availability. The complete nucleotide sequence of this SARS-CoV-2 isolate (SARS-CoV-2/human/BGD/NIB_01/2020) has been deposited in GenBank under the accession number MT509958.




FIG 1 Phylogenetic analysis of the SARS-CoV-2/human/BGD/NIB_01/2020 isolate. Nucleotide alignment and the GTR+CAT nucleotide evolution model was applied to construct the tree. The tree was visualized using i-TOL. Here, the x axis represents the tree scale. A scale bar with a 0.0001 value is given on the top. The genome (labeled in red) shares a common ancestor with some isolates from the United States.

ACKNOWLEDGMENTS

We are grateful to the Ministry of Science and Technology for its extensive support during this research work.

This study was funded by the National Institute of Biotechnology, Ministry of Science and Technology, government of the People's Republic of Bangladesh.

REFERENCES

- Besser J, Carleton HA, Gerner-Smidt P, Lindsey RL, Trees E. 2018. Next-generation sequencing technologies and their application to the study and control of bacterial infections. *Clin Microbiol Infect* 24:335–341. <https://doi.org/10.1016/j.cmi.2017.10.013>.
- Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet* 16:276–277. [https://doi.org/10.1016/S0168-9525\(00\)02024-2](https://doi.org/10.1016/S0168-9525(00)02024-2).
- Zhang Z, Schwartz S, Wagner L, Miller W. 2000. A greedy algorithm for aligning DNA sequences. *J Comput Biol* 7:203–214. <https://doi.org/10.1089/10665270050081478>.
- Kazutaka K, John R, Kazunori DY. 2019. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 20:1160–1166. <https://doi.org/10.1093/bib/bbx108>.
- Price MN, Dehal PS, Arkin AP. 2010. FastTree 2: approximately maximum-likelihood trees for large alignments. *PLoS One* 5:e9490. <https://doi.org/10.1371/journal.pone.0009490>.
- Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res* 46:W537–W544. <https://doi.org/10.1093/nar/gky379>.
- Letunic I, Bork P. 2019. Interactive Tree of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res* 47:W256–W259. <https://doi.org/10.1093/nar/gkz239>.